

Strangeness Production Experiments at Jefferson Lab*

Reinhard A. Schumacher

*Department of Physics,
Carnegie Mellon University,
Pittsburgh, PA 15213, USA
E-mail: schumacher@cmu.edu*

Experimental results for photo- and electro-production of open strangeness from the Thomas Jefferson National Accelerator Facility are discussed. The results are from work completed by mid-2003 on elementary KY production, nuclear targets, and the exotic Θ^+ state. It is shown how the increases in intensity and precision of JLab experiments over earlier work have allowed new phenomena to become measurable.

I. INTRODUCTION

Electro- and photo-production of open strangeness is interesting because it involves a well-understood probe – the photon – interacting with a nucleon or nuclear target to make an $s\bar{s}$ quark pair that is not part of the target’s valence structure. This makes reactions in which strange particles are produced attractive to look for quark-level dynamics in the production mechanism. Two examples of possible quark-level dynamics are shown in this report. Where direct signatures of quark degrees of freedom are not evident, meson and baryon descriptions of the strangeness production mechanism are sensitive to the inclusion of non-strange excited states of nucleons that decay to strange particles in the final state. For example, excited non-strange baryons which are “hidden” when searching in pionic final states may be “revealed” by looking in final states with strangeness, simply because couplings to pion- and kaon-containing final states will, in general, be different. In the area of strangeness production off nuclei, information about the hyperon-nucleon interaction can be gleaned from final state interactions with a deuteron target; a first result in this field has been obtained. The creation of hypernuclear states has been demonstrated at Jefferson Lab for the first time, opening a new field of research for the future. The final topic in this report is the recent evidence from Jefferson Lab of a strangeness +1 baryon dubbed the Θ^+ . At Jefferson Lab, this state has been seen with CLAS in an exclusive reaction on the deuteron, as well as in a different final state off the proton.

This report gives an overview of the experimental results that have been obtained to date at Jefferson Lab in these fields. It also mentions the experiments that are still under

* Submitted to the Proceedings of “SENDAI03: Electrophoto-production of Strangeness on Nucleons and Nuclei”, Tohoku University, Sendai, Japan, June 2003. To be published by World Scientific.

analysis or that have yet to take data. The sections we will use for the discussion are:

- Elementary production — on the proton
- Nuclear electroproduction — on the deuteron and heavier nuclei
- The Θ^+ state — first results from CLAS

II. ELEMENTARY PRODUCTION ON THE PROTON

The elementary photo- and electro-production of hyperons in the resonance region is used mainly as a test of production mechanisms through resonant baryon formation and kaon exchange. Couplings of N^* 's and Δ 's to KY final states differ from couplings to pions and nucleons but are not well known. Thus, one can expect to learn about how the couplings compare to expectations (from flavor SU(3), for instance), and possibly also to “see” resonance contributions that are hidden in the case of pionic final states.




FIG. 1: Missing mass spectrum for photoproduction of strangeness in CLAS, using a tagged bremsstrahlung beam with 2.4 GeV endpoint energy.

A. Photoproduction

CLAS [1] has measured [2] the cross sections and recoil polarizations for the reactions $\gamma + p \rightarrow K^+ + \Lambda$ and $\gamma + p \rightarrow K^+ + \Sigma^0$ from threshold up to 2.3 GeV. The mass resolution for the detected kaons was $\sigma = 6.1$ MeV, as determined from time of flight. This is shown in Fig. 1. One of the main results from this work is the energy dependence of the $K^+\Lambda$ cross sections shown in Fig. 2. The panels show how, as a function of the s -channel energy, W , the

cross section evolves at very forward (top), forward (middle), and backward (bottom) center-of-mass angles. At the extreme angles there are peaks near threshold and near 1.9 GeV, while in the middle-forward region the dip between the peaks is filled in. The higher-energy peaks are significant since they confirm hints of this structure published by the SAPHIR collaboration [7]. They indicate that there are at least two structures in the mass range near 1.9 GeV which couple strongly to $K^+\Lambda$, since the shape of the peaks and their position change somewhat with angle. Three calculations are shown for reference. To mention just one feature, the MAID model [3] is based on the hadrodynamic model of Mart *et al.* [4], and generates the higher energy peak with a $D_{13}(1895)$ state. Clearly this is not enough to describe the data. These data will now allow more detailed modeling of the production of $K^+\Lambda$ to be constructed.




FIG. 2: Energy dependence of the Λ cross section at the most forward angle measured (top), and at intermediate forward and backward angles (middle, bottom); from McNabb *et al.* [2]. The curves are for effective Lagrangian calculations computed by Kaon-MAID [3, 4] (solid), the related model of Janssen *et al.* [5] (dashed), and a Regge-model calculation of Guidal *et al.* [6] (dot-dashed). Data from SAPHIR [7] (open triangles) are also shown.

The hyperon recoil polarization has also been measured, as can be seen in the un-

published portion of Ref [2]. CLAS also has data under analysis of photoproduction using linearly and circularly polarized photon beams. The observables Σ (linear polarized beam asymmetry) [8], C'_x and C'_z (beam-recoil double polarizations) [9] will eventually be available, measured over the same kinematic range as the results mentioned above.

B. Electroproduction

Moving away from the kinematic point at $Q^2 = 0$, electroproduction of strangeness introduces two new ingredients into the discussion: the longitudinal coupling of the photons in the initial state, and the electromagnetic and hadronic form factors of the exchanged particles. New observables include the longitudinal-transverse and transverse-transverse interference terms in the cross section. These have been measured by CLAS for the first time over a broad kinematic range. The expression for the unpolarized electroproduction cross section is, after dividing out the flux of virtual photons,

$$d\sigma/d\Omega_K = \sigma_U + \epsilon\sigma_{TT} \cos(2\phi) + \sqrt{2\epsilon(\epsilon+1)}\sigma_{LT} \cos(\phi), \quad (1)$$

where $\sigma_U = \sigma_T + \epsilon\sigma_L$ is the unseparated sum of the transverse and longitudinal cross sections, ϵ is the photon polarization parameter determined by the electron scattering kinematics, and ϕ is the azimuthal angle between the electron scattering and the hadronic reaction planes. In CLAS it is possible to detect the scattered electrons together with the produced kaons over a broad range of kinematics to separate out the ϕ angle dependence of the reaction, and thus determine σ_U , σ_{LT} , and σ_{TT} . A sample result [10] is shown in Fig. 3, where three components of the cross section are plotted as a function of W for both hyperons. The data and model calculations have been averaged over $0.5 < Q^2 < 0.9 \text{ GeV}^2$, and also averaged over the ‘backward’ kaon center-of-mass angles of $-0.67 < \cos(\theta_K) < -0.33$.

A major reason the cross section components for Λ and Σ^0 are different in Fig. 3 is due to the isospin selectivity, *i.e.* the $K^+\Lambda$ final state couples only to N^* ’s, while $K^+\Sigma^0$ ’s couple to both N^* ’s and Δ ’s. One sees that σ_U in the Λ cross section has two peaks, as was seen in the photoproduction data, thus confirming the strong contribution of a resonance-like structure near 1.9 GeV as seen in the photoproduction data. In the Σ^0 case there is only a broad peak without clear structure. In the interference cross sections the most prominent feature at these kinematics is a broad negative dip in σ_{TT} for the Σ^0 case. One notes immediately that none of the models (the same models cited previously) are able to consistently describe these results. However, there are several well-established Δ^* resonances near 1.9 GeV which can couple to $K^+\Sigma^0$. This means that more work needs to be done to model resonance content and the Q^2 dependence of the reaction mechanisms.

The circular polarization of virtual photons can be transferred to the produced Λ ’s in electroproduction, and this process has been studied at CLAS [11] using a $\sim 70\%$ polarized electron beam. The “beam-recoil” double polarization thus measured is again amenable to




FIG. 3: Energy dependence of the Λ (left) and Σ^0 (right) cross sections for $\sigma_U = \sigma_T + \epsilon\sigma_L$ (top), σ_{TT} (middle), and σ_{LT} (bottom). The curves are for effective Lagrangian calculations computed by Kaon-MAID [3] (solid), the similar model of Janssen *et al.* [5] (dot-dashed), and a Regge-model calculation of Guidal *et al.* [6] (dashed).

comparison with a variety of reaction models. The two components of Λ polarization in the $K\Lambda$ plane can be determined by measuring the beam-helicity-dependent yield asymmetry of protons from the decay of the Λ 's. The main result is that the polarization of the Λ 's is maximal in the direction of the virtual photon, and minimal transverse to this direction. Fig. 4 shows the data. This phenomenon can be qualitatively understood [11] at the quark level in terms of the $s\bar{s}$ quark pair being formed predominantly in the anti-aligned state, with the produced s -quark carrying the Λ spin, polarized along the photon helicity direction. It may also be a manifestation of the VDM concept that the photon is first converted into an aligned virtual $s\bar{s}$ pair in the form of a virtual ϕ meson followed by quark rearrangement.

For the reactions $p(e, e'K^+)Y$ it has also been possible to make a separation of the pieces of σ_U into transverse and longitudinal cross section using the traditional Rosenbluth method. This has been done in Hall C at Jefferson Lab using the “Short Orbit Spectrometer” to detect kaons in coincidence with the “High Momentum Spectrometer” to detect scattered electrons [12]. The acceptance of these magnetic spectrometers led to a very narrow set of kinematics at which the separation was made: specifically, the kaon was along the direction




FIG. 4: Polarization transfer to the recoiling hyperon in the reaction $\bar{e} + p \rightarrow e' + K^+ + \bar{\Lambda}$. Top panel shows the large Λ polarization along the photon direction, bottom panel shows the vanishing polarization transverse to this direction. From Carman *et al.* [11].

of the virtual photon, *i.e.* $\theta_{\gamma K} = 0^\circ$, and for $W = 1.84$ GeV. Fig. 5 shows the resulting cross section components and their ratio, for Λ 's and $\Sigma^{0\prime}$ s, as a function of Q^2 . These are the most precise separations ever measured for these hyperons. They show that the ratio σ_L/σ_T is about one half at these kinematics up to $Q^2 = 2$ GeV².

The calculations shown for comparison exhibit, at best, only fair agreement with the data. The work of Williams, Ji, and Cotanch [13] is perhaps the better of the two, at least for the ratio of σ_L/σ_T for the Λ . The authors of the paper [12] surmise that the baryon resonance content of the models, as well as the modeling of the hadronic form factors must be improved in order to successfully reproduce these results. This is the same conclusion that can be reached from examination of the CLAS data shown above. Generally, all of these new high-precision data for photo- and electro-production of strangeness on the proton call out for a renewed effort at understanding baryon and meson exchange structure of these reactions.

It would be very interesting to measure the charge form factor of the K^+ meson in order to compare it with the form factor of the π^+ . The same Hall C data set used in the analysis of the Mohring *et al.* result discussed above has been used by the E93-018 Collaboration [14] to extrapolate the t -dependence of σ_L to the kaon pole (the Chew-Low method). A single unpublished form factor value $Q^2 = 1.0$ GeV² has been obtained. Further analysis of data for values at $Q^2 = 0.5$ and 2.0 GeV², taken in Halls C and A, is in progress in order to determine the trend of this form factor.




FIG. 5: Results from Mohring *et al.* [12] for Rosenbluth separation of σ_L (a), σ_T (b), and their ratio (c). The data are compared to calculations of Refs. [13] and [4]. Results for the Λ are at left and Σ^0 at right.

The decay of the electroproduced excited hyperon $\Lambda(1520) \rightarrow K^- p$ was studied at CLAS [15]. The angular distribution in the t -channel helicity frame gives information about whether the t -channel exchange particle had spin $J = 0$ (such as the K^-) or had J greater than 0 (such as the K^{*-}). This is interesting because in photoproduction, at $Q^2 = 0$, there is a clear preference for $\Lambda(1520)$ formation via K^{0*} exchange. At $Q^2 > 0.9 \text{ GeV}^2$, the CLAS experiment showed that $J = 0$ exchange provides about 60% of the formation channel. These results are striking, but have not yet lead to renewed theoretical activity on this reaction mechanism.

III. PRODUCTION FROM NUCLEI

The Jefferson Lab data on hyperon electroproduction on the deuteron and heavier nuclei have recently started to appear in archival journals. Other results have been shown at conferences only. Below we give an overview of the most interesting new results. They come from Hall C work, but there is also program getting underway in Hall A.

A. The YN Interaction

Quasi-free kaon production on the deuteron, $d(e, e' K^+) YN$, has been measured by the E91-016 Collaboration in Hall C [16]. The significance of this work is two-fold: it gives unique access to one of the elementary production reactions, $n(e, e' K^+) \Sigma^-$, and secondly,

on one side of the quasi-free peak there is a kinematic regime where the hyperon and the nucleon have nearly the same momentum, and therefore interact via the low energy YN interaction as a final state interaction (FSI).

Figure 6 shows the reaction at $Q^2 = 0.38 \text{ GeV}^2$ and a beam energy of 3.2 GeV. The curves are Monte Carlo calculations which do not include FSI. The contribution from the neutron to Σ^- production is substantial. On the left-hand edge one sees the deficit in the prediction of the quasi-free Monte Carlo model which must be filled in by FSI. Several YN models have been used to compute the final state interaction in the simulations (not shown here), each with a characteristic S-wave scattering length and effective range; some sensitivity has been shown, favoring the “Verma” and “Jülich A” potentials and not favoring “Jülich B”.




FIG. 6: Reconstructed missing mass from $d(e, e' K^+)$ from Ref [16]. The curves are Monte Carlo simulations which do not have YN final state interactions turned on. FSI effects are seen as excess counts near the Λd threshold.

B. Hypernuclear Electroproduction

A remarkable achievement of the Jefferson Lab program has been the observation of electromagnetically produced hypernuclear states in light nuclei. Experiments using two different approaches have each succeeded in finding evidence for such states.

Using the standard Hall C spectrometers, the E91-016 Collaboration [17] measured the reactions $A(e, e' K^+) YX$ for $^1H, ^2H, ^3He, ^4He, C$, and Al targets, at $Q^2 = 0.35 \text{ GeV}^2$ and for the lab angle between the virtual photon and the kaon at $0^\circ, 6^\circ$, and 12° . The most compelling result was for the 4He target, shown in Fig. 7 which showed clear bound

state peaks at the end of the quasi-free production spectrum for all three scattering angles. These peaks correspond to the formation of the well-known hypernucleus ${}^4_{\Lambda}H$. The cross section was estimated to be about 20 nb/sr. The data in the figure shows no evidence of Σ hypernuclear production (dashed lines high in quasi-free spectrum).




FIG. 7: Missing mass distributions for ${}^4He(e, e'K^+)$ showing quasi-free Lambda production and the formation of the bound state ${}^4_{\Lambda}H$, from Dohrmann *et al.* [17].

The limitation of the experimental technique used to produce this result is that the missing-mass resolution is insufficient to extend these measurements to heavier hypernuclei where many excited states are present. Another method which achieves resolution of under 1 MeV would be required to make electromagnetic hypernuclear production truly interesting as a research tool for hypernuclear spectroscopy.

This other method was pioneered by the E89-009 collaboration [18]. They used the Hall C SOS spectrometer to detect kaons, but used a dedicated dipole “Enge splitpole” spectrometer for the electrons. In addition, a beam-splitter magnet was used to allow the electron angle to be pushed all the way to zero degrees, thus decreasing Q^2 and maximizing the rate of good events. Also, great care was taken to minimize straggling effects in materials. The result was obtained [18] for the reaction ${}^{12}C(e, e'K^+){}^{12}_{\Lambda}B$, seen in Fig. 8, which showed peaks corresponding to bound Λ 's in the S and in the P shells.

The experiment achieved a missing mass resolution of 0.9 MeV FWHM, which can be compared with about 1.5 MeV as the best value achieved using the (π^+, K^+) reaction for




FIG. 8: Missing mass distributions for $^{12}\text{C}(e, e'K^+)^{12}_{\Lambda}\text{B}$ showing peaks corresponding to the ground state (Λ in the S shell), and P-shell excited state (Λ in the P shell). The curve shows predictions for some core-excited states in which the host ^{11}B nucleus is not in its ground state. From Ref [18].

hypernuclear spectroscopy. The experiment was limited by statistics and resolution, so that little additional quantitative information has emerged. The experiment demonstrated that spectroscopic resolution is achievable, but to turn the method into a research tool a better instrument is needed. A new kaon spectrometer has been built [19] that should achieve 0.4 MeV resolution and permit count rates of over several 100 per day for the $^{12}_{\Lambda}\text{B}$ ground state. It will be installed at Jefferson Lab in mid-2004.

In Hall A at Jefferson Lab the E94-107 Collaboration [20] is also planning to start hypernuclear spectroscopy in late 2003. They will use the existing HRS spectrometers, augmented by a septum-magnet arrangement which will allow the electron spectrometer to reach 6° in the lab and by a RICH detector for more effective K/π separation.

IV. THE Θ^+ STATE

In the naive quark model, baryons come only in colorless 3-quark flavor combinations. However, no fundamental rule forbids combinations of more quarks as long as they are also colorless. Hence, for decades there have been searches for other multi-quark structures that would, if they exist, be important ingredients of hadronic physics. No such unusual states had been convincingly identified however, so the Fall 2002 announcement by the

LEPS group at SPring-8 [21] in Japan of a narrow pentaquark state near 1540 MeV that decays into K^+n caused renewed excitement. This state is manifestly “exotic” since the K^+ contains an anti- s quark, which cannot be an ingredient in an ordinary baryon. A large number of theoretical papers have appeared in recent months, as well as at least 4 experimental confirmations [21, 22, 23, 24], showing that this topic has become very hot. Spring-8’s LEPS group was inspired by the theoretical prediction of Diakonov, Petrov, and Polyakov [25] who predicted a narrow $uudd\bar{s}$ pentaquark state within the framework of a chiral soliton model, in which it appears as one member of an anti-decuplet of 5-quark states. The observation of a state with some of the predicted features is not yet final proof, of course, that this model is the optimal explanation.

The CLAS Collaboration at Jefferson Lab has recently reported [23] results from the exclusive reaction $\gamma d \rightarrow K^+ K^- pn$ in which the neutron is reconstructed via the missing mass off the three detected charged particles. Two of many possible diagrams through which the Θ^+ might be formed are shown in Fig. 9. The “two-step” nature of the production mechanism has the experimental benefit that the K^- ’s that are typically very forward peaked rescatter to larger lab angles, and therefore are more inside the CLAS acceptance.




FIG. 9: Diagrams leading to the production and decay of a Θ^+ from exclusive reactions on the deuteron.

The Θ^+ signal was sought in the invariant mass of the K^+n decay channel, and the final result is shown in Fig. 10. The signal is deduced to have about “5 sigma” significance, using a smooth (polynomial) background. Various background shapes have been tested, with similar significances. Production of excited hyperons is present in the same final state, and forms background under the peak in the final spectrum; the $\Lambda(1520)$ events were removed explicitly, however, with a cut on the K^-p mass distribution. Full Monte Carlo modeling of the background has not been completed, although preliminary tests have suggested that the background has a large contribution from non-resonant KK production off the bound nucleon.




FIG. 10: Invariant mass of the K^+n system, which has strangeness $S = +1$, showing a sharp peak at the mass of 1.548 (GeV/c)^2 . The fit (solid line) to the peak on top of the smooth background (dashed line) gives a statistical significance of about 5σ . The dashed histogram shows events removed by the hyperon cut, and the dotted histogram shows the scaled background from non-pKK events.

Immediately one wonders whether the Θ^+ state could also be reached in photo-production off the proton directly. At CLAS there is a search at this time for the reaction $\gamma + p \rightarrow \bar{K}^0 + \Theta^+ \rightarrow \pi^+\pi^-K^+n$. A positive result has been reported recently from SAPHIR [24], in the same channel. Interestingly, CLAS does have evidence for the Θ^+ in the reaction $\gamma + p \rightarrow K^- + \pi^+ + \Theta^+ \rightarrow K^- + \pi^+ + K^+ + n$. (The result was shown in the oral presentation but is not included here since the data are still in flux.) In conclusion, the study of this new exotic baryonic state is very intense at this time, with positive sighting of a narrow Θ^+ by at least 4 experimental groups, and with a plethora of theoretical interpretation underway.

V. OTHER WORKS IN PROGRESS

Additional work in progress on strangeness production at Jefferson Lab has not been mentioned so far due to time and space constraints. In the future we hope to present results on the following topics: (i) the radiative decay of the $\Sigma(1385)$, which tests quark-model wavefunctions [26], (ii) cross sections [27] for $\gamma p \rightarrow \bar{K}^0\Sigma^+$ (iii) the beam-recoil double polarization observables C'_x and C'_z in $p(\gamma, K^+)\Lambda$ [9], (iv) cross sections for K^+ photoproduction off the deuteron [28], (v) results for the “fifth” structure function in electroproduction [29], (vi) photoproduction cross sections for the excited hyperons [30], (vii) line-shape analysis for

photoproduction of the $\Lambda(1405)$ decaying to different $\Sigma\pi$ charge states, as a test of its structure [31]; (viii) photoproduction of $S = -2$ Cascade (Ξ^-) states [32].

VI. SUMMARY

The disparate experimental results presented here make it difficult to summarize this report in a few words or a single central new concept. The main observations are that (a) interesting new baryon resonance structure has been seen in both photo- and electroproduction of strangeness on the nucleon; (b) electroproduction off the deuteron has shown that it is possible to test models of the YN final state interaction; (c) hypernuclear production from ^3He and ^4He and from ^{12}C shows the feasibility of using electromagnetic probes to study production and structure of hypernuclei; and (d) an exotic new baryon, the Θ^+ , has been confirmed to exist at Jefferson Lab, a discovery which is engendering a great deal of new theoretical and experimental work.

Acknowledgments

The author wishes to thank numerous colleagues who made their results available for this talk and paper. They include K. Baker, S. Barrow, E. Beise, D. Carman, F. Dohrmann, K. Hicks, J. Melone, R. Feuerbach, V. Koubarovski, K. Livingston, P. Markowitz, J. McNabb, M. Mestayer, G. Niculescu, J. Price, J. Reinhold, S. Stepanyan, and L. Tang. They and other colleagues offered advice and suggestions which are gratefully acknowledged. This work is supported by DOE contract DE-FG02-87ER40315.

-
- [1] B. Mecking *et al.*, Nucl. Instrum. and Methods **A503**, 513 (2003), and references therein.
 - [2] J.W.C. McNabb *et al* *nucl-ex/0305028*; submitted to Phys. Rev. Lett.; W. J. C. McNabb Ph.D. Thesis, Carnegie Mellon University (2002) (unpublished).
 - [3] T. Mart, C. Bennhold, H. Haberzettl, and L. Tiator, “KaonMAID 2000” at www.kph.uni-mainz.de/MAID/kaon/kaonmaid.html.
 - [4] T. Mart and C. Bennhold, Phys. Rev. **C61**, 012201 (2000); C. Bennhold, H. Haberzettl, and T. Mart, *nucl-th/9909022* and Proceedings of the 2nd Int’l Conf. on Perspectives in Hadronic Physics, Trieste, S. Boffi, ed., World Scientific, (1999).
 - [5] S. Janssen, J. Ryckebusch, D. Debruyne, and T. Van Cauteren, Phys. Rev **C65**, 015201 (2001); S. Janssen *et al.*, Eur. Phys. J. **A 11**, 105 (2001); curves via private communication.
 - [6] M. Guidal, J.-M. Laget, and M. Vanderhaeghen, Phys Rev. **C61**, 025204 (2000); M. Vanderhaeghen, M. Guidal, and J.-M. Laget, Phys. Rev. **C57**, 1454 (1998).
 - [7] M. Q. Tran *et al.*, Phys. Lett. **B445**, 20 (1998); M. Bockhorst *et al.*, Z. Phys. **C63**, 37 (1994).

- [8] CLAS Experiment 98-109, P. Cole, spokesperson, and K. Livingston and J. Melone, (private communication).
- [9] CLAS Experiment 89-004, R. Bradford, PhD thesis.
- [10] R. Feuerbach *et al.*, to be published. Ph.D. Thesis, Carnegie Mellon University (2002) (unpublished).
- [11] D. S. Carman *et al.*, *Phys. Rev. Lett.* **90**, 131804 (2003); M. Mestayer, these proceedings.
- [12] R. M. Moring *et al.*, *Phys. Rev.* **C67**, 055205 (2003). See also the earlier paper for the same experiment: G. Niculescu *et al.*, *Phys. Rev. Lett.* **81**, 1805 (1998).
- [13] R.A.Williams, C.-R. Ji, and S. R. Cotanch *Phys. Rev.* **C46**, 1617 (1992).
- [14] O. K. Baker for the E93-018 collaboration (private communication).
- [15] S. P. Barrow *et al.*, *Phys. Rev.* **C64**, 044601 (2001).
- [16] J. Reinhold for the E91-016 collaboration, Proceedings of Baryons 2002, C. Carlson and B. Mecking, Eds., World Scientific, 589, (2002), and private communication.
- [17] F. Dohrmann for the E91-016 collaboration, Proceedings of Baryons 2002, C. Carlson and B. Mecking, Eds., World Scientific, 585, (2002), and private communication.
- [18] T. Miyoshi *et al.*, *Phys. Rev. Lett.* **90**, 232502 (2003).
- [19] Y. Fujii *et al.*, *Nucl. Phys.* **A721**, 1079c (2003).
- [20] P. Markowitz for the E94-107 collaboration (private communication).
- [21] T. Nakano *et al.*, *Phys. Rev. Lett.* **91**, 092001 (2003).
- [22] V. V. Barmin *et al.*, hep-ex/0304040, (2003); accepted by *Yad. Phys.*
- [23] S. Stepanyan *et al.*, hep-ex/0307018, (2003); submitted to *Phys. Rev. Lett.*
- [24] J. Barth *et al.*, hep-ex/0307083, (2003).
- [25] D. Diakonov, V. Petrov, and M. Polyakov, *Z. Phys.* **A359**, 305 (1997).
- [26] CLAS Experiment 89-024, G. Mutchler, spokesman.
- [27] CLAS Experiment 89-004, B. Carnahan, Ph.D. Thesis, Catholic Univ. of America.
- [28] CLAS Experiment 89-045, B. Mecking, spokesman, I. Niculescu, contactperson.
- [29] CLAS Experiment 99-006, D. Carman and B. Raue, spokesmen.
- [30] CLAS Experiment 89-004, H. Juengst, contactperson.
- [31] CLAS Experiment 89-004, R. Schumacher, spokesman.
- [32] CLAS Approved Analysis, J. Price, spokesman.